



P: (843) 353-3460 | F: (843) 353-3461

DISTAL BICEPS REPAIR Postoperative Instructions

I. Diet

- Begin with clear liquids and light foods (Jell-O, soups, etc.)
- Progress to your normal diet if you are not nauseated.

II. Wound Care

- Maintain your operative dressing, loosen bandage if swelling of the hand occurs.
- It is normal for the elbow to bleed and swell following surgery. If blood soaks through the bandage, do not become alarmed, reinforce with additional dressing.
- To avoid infection, keep surgical incisions clean and dry – you may shower by placing a plastic covering over the surgical site beginning the day after surgery. NO immersion of the arm.
- Elbow will be immobilized in a splint/sling following surgery do not remove until post op appointment 7-10 days following surgery. You will be transitioned to an elbow brace at this time.
- You may begin showering and getting your wound site wet once the splint is removed.

III. Medications

- Local anesthetics are injected into the wound at the time of surgery. This will wear off within 8-12 hours and it is not uncommon for patients to encounter more pain on the first or second day after surgery when swelling peaks.
- Most patients will require some narcotic pain medication for a short period of time – this can be taken as per directions on the bottle.
 - Primary Medication = (Oxycodone)
 - Take 1 – 2 tablets every 4 – 6 hours as needed
 - Max of 12 pills per day
 - Plan on using it for 2 to 5 days, depending on level of pain
 - Ondansetron (Zofran) take as needed for nausea
 - Docusate sodium (Colace) as needed for constipation
- Meloxicam two times a day for 2 weeks
- Tylenol 1000mg three times a day for two weeks
- Common side effects of the pain medication are nausea, drowsiness, and constipation. To decrease the side effects take the medication with food. If constipation occurs, consider taking an over the counter laxative.

- If you are having problems with nausea and vomiting, contact the office to possibly have your medications changed.
- Do not drive a car or operate machinery while taking the narcotic medication.

IV. Activity

- When sleeping or resting, inclined positions (i.e.: reclining chair) and a pillow under the forearm for support may provide better comfort.
- Do not engage in activities which increase pain/swelling. Unless otherwise instructed the arm should remain in the sling at all times.
- Avoid long periods of sitting or long distance traveling for 2 weeks.
- NO driving until instructed otherwise by physician
- May return to sedentary work ONLY or school 3-4 days after surgery, if pain is tolerable
- Formal physical therapy typically begins 7-10 days postoperatively. Please call your physical therapy location of choice to schedule appointments within 2-3 days after surgery.

V. Immobilizer (If Prescribed)

- You will be placed in a posterior mold splint where you cannot straighten your elbow for 1-2 weeks following surgery. You are to wear sling at all times while the splint is in place.
- You will be transitioned to an elbow brace at your post-operative appointment.

VI. Ice Therapy

- Icing is very important in the initial post-operative period and should begin immediately after surgery.
- Ice packs 20 minutes every 2 hours daily until your first post-operative visit. Care should be taken with icing to avoid frostbite to the skin.

VII. Exercise

- Begin finger flexion and extension on the first post-operative day to help decrease swelling.
- Formal physical therapy (PT) and occupational therapy (OT) typically begins 7-10 days after surgery. Please contact your therapy location of choice to schedule an appointment 2-3 days after surgery.
- If you need a referral faxed to an outside location, please contact our office.

VIII. Emergencies

- Contact Dr. Gehrman or his clinical assistant at (843) 353-3460 if any of the following are present:
 - Painful swelling or numbness
 - Unrelenting pain

- Fever (over 101°F - it is normal to have a low grade fever for the first day or two following surgery) or chills
- Redness around incisions
- Color change in wrist or hand
- Continuous drainage or bleeding from incision (a small amount of drainage is expected)
- Difficulty breathing
- Excessive nausea/vomiting
- **If you have an emergency after office hours or on the weekend, contact the same office number 843-353-3460 and you will be connected to our page service – the will contact Dr. Gehrman or one of his colleagues if he is unavailable. Do NOT call the hospital or surgical center.
- **If you have an emergency that requires immediate attention, proceed to the nearest emergency room.

IX. Follow Up Care + Questions

- A member of Dr. Gehrman's team will call you 24 to 48 hours after surgery to address any questions or concerns. If you have not been contacted within 48 hours of surgery, please call the office at 843-353-3460.
- Typically the first post-operative appointment following surgery is 10-14 days following surgery.
- If you do not already have a postoperative appointment scheduled, please contact the office during normal office hours (843-353-3460) and ask for appointment scheduling.
- The first post-operative appointment will be with myself or one of the Physician Assistants. They will assess the wound, go over post-operative protocol, and answer any questions you may have regarding the procedure.

DISTAL BICEPS REPAIR REHABILITATION PROTOCOL

Name: _____ **Date:** _____

Diagnosis: _____

Date of Surgery _____

Frequency: 1 2 3 4 times/week

Duration: 1 2 3 4 5 6 Weeks

Weeks 0-4:

- Elbow is immobilized in the Bledsoe Brace at 75 degrees flexion with wrist free.
- Dressing changed at 7-10 days after surgery
- Neck strap / sling to be used for the first 2-5 days. Then, may begin AROM of shoulder

Weeks 4-6:

- Brace adjusted to 30 degrees extension, may begin active assisted flexion to 90 degrees in brace

Weeks 6-8:

- Discontinue the use of the Bledsoe Brace
- Shoulder and elbow ROM, PROM-AAROM-AROM, advance as tolerated • Begin muscle strengthening exercises for wrist and forearm

Months 2-3:

- May begin elbow strengthening

